



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

MESTRADO INTEGRADO EM MEDICINA

2015/2016

Ana Filipa Rodrigues Félix
Main Causes of Sudden Deaths in
Pregnancy

março, 2016

FMUP



FACULDADE DE MEDICINA
UNIVERSIDADE DO PORTO

Ana Filipa Rodrigues Félix
Main Causes of Sudden Deaths in
Pregnancy

Mestrado Integrado em Medicina

Área: Medicina Legal

Tipologia: Monografia

Trabalho efetuado sob a Orientação de:
Doutor Agostinho Santos

Trabalho organizado de acordo com as normas da revista:
Journal of Forensic and Legal Medicine

março, 2016

FMUP

Eu, Ana Filipa Rodrigues Félix, abaixo assinado, nº mecanográfico, 201001192 estudante do 6º ano do Ciclo de Estudos Integrado em Medicina, na Faculdade de Medicina da Universidade do Porto, declaro ter atuado com absoluta integridade na elaboração deste projeto de opção.

Neste sentido, confirmo que **NÃO** incorri em plágio (ato pelo qual um indivíduo, mesmo por omissão, assume a autoria de um determinado trabalho intelectual, ou partes dele). Mais declaro que todas as frases que retirei de trabalhos anteriores pertencentes a outros autores, foram referenciadas, ou redigidas com novas palavras, tendo colocado, neste caso, a citação da fonte bibliográfica.

Faculdade de Medicina da Universidade do Porto, 15/02/2016

Assinatura conforme cartão de identificação:

Ana Filipa Félix

NOME

Ana Filipa Rodrigues Félix

NÚMERO DE ESTUDANTE

201001192

DATA DE CONCLUSÃO

DESIGNAÇÃO DA ÁREA DO PROJECTO

Medicina legal

TÍTULO DISSERTAÇÃO/MONOGRAFIA (riscar o que não interessa)

Main Causes of Sudden Deaths in Pregnancy

ORIENTADOR

Doutor Agostinho Santos

COORDENADOR (se aplicável)

ASSINALE APENAS UMA DAS OPÇÕES:

É AUTORIZADA A REPRODUÇÃO INTEGRAL DESTA TRABALHO APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
É AUTORIZADA A REPRODUÇÃO PARCIAL DESTA TRABALHO (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) APENAS PARA EFEITOS DE INVESTIGAÇÃO, MEDIANTE DECLARAÇÃO ESCRITA DO INTERESSADO, QUE A TAL SE COMPROMETE.	<input type="checkbox"/>
DE ACORDO COM A LEGISLAÇÃO EM VIGOR, (INDICAR, CASO TAL SEJA NECESSÁRIO, Nº MÁXIMO DE PÁGINAS, ILUSTRAÇÕES, GRÁFICOS, ETC.) NÃO É PERMITIDA A REPRODUÇÃO DE QUALQUER PARTE DESTA TRABALHO.	<input checked="" type="checkbox"/>

Faculdade de Medicina da Universidade do Porto, 23/3/2016

Assinatura conforme cartão de identificação: Ana Filipa Felix

“À minha mãe. “

Main Causes of Sudden Deaths in Pregnancy

Ana Félix MD (filipa_felix5@hotmail.com)

Faculdade de Medicina do Porto, Porto, Portugal

Agostinho Santos MD, PhD,

Faculdade de Medicina do Porto, Porto, Portugal

Abstract

Background The pregnancies represent a big physiologic change, although in the majority of the cases this alteration are easily exceed, sometimes, associated or not with previous pathologies can cause important complication to the pregnant women. The low numbers of registered cases hinder then a more specific study on the subject. The aim of this review is explore the main causes of sudden death directly and indirectly related with pregnancy.

Materials and methods Using article databases such as PubMed and books of obstetrics and legal medicine and organizations of the last ten years, we collected, and integrated information about the etiology of maternal sudden death including definition, physiopathology, epidemiology and risk factors, clinic, diagnosis, and autopsy findings.

Results The last results from Directorate-General of Health in 2013 features a maternal mortality of 8 deaths per 100000 live births. Between 2001 and 2007, in Portugal, 21.7% of the total number of maternal deaths were of unknown cause.

The main direct causes are hemorrhage/coagulation disorder, hypertensive disease of pregnancy, infection/sepsis and embolism. Pathologies such as heart and neurologic diseases, cancer disease and infection are the most important indirect causes.

Discussion Sudden death in pregnancy has a big impact in the society but actually in Europe it has been little explored.

Conclusion In order to decrease maternal death is essential the establishment of national and international reports with updated statistics and awareness of physicians to the importance of the implementation of the screening measures and treatment filed for each stage of pregnancy.

Key words: *pregnancy, sudden death, direct causes, indirect causes, autopsy*

Introduction

The World Health Organization (WHO) defines Maternal Mortality as the deaths of women during pregnancy, childbirth or in the 42 days of puerperium, independent of the duration and the local of the pregnancy. The concept of Maternal Mortality includes all the causes that can be associated or aggregated with pregnancy or its management. According to this definition, ectopic pregnancy and abortion are included in the causes, but the incidental deaths are excluded.^[3] It is one of the most important indexes of the health care service of a country.^[4]

The last data publish by WHO, in 2015, indicated that Maternal Mortalities are 12 and 239 per 100000 live births in developed and developing countries respectively. The first one has a lifetime risk of maternal death of 1/4900 comparing with the 1/150 in developing countries.(table 4)^[2]

The absolute numbers have been decreasing in 44% in the last 25 years, with the technological evolution, better access to health care and the reduction of pregnancies in extreme ages.^[2]

United Nations created the Millennium Development Goals (MDGs), which include maternal death surveillance and response (MDSR) in order to improve the precision of the notification of the deaths. The aim of MDSR was to decrease maternal mortality in 75% at 2015. The new aim is to get less than 30 deaths per 100000 pregnancies until 2030.^[1] Primary prevention is, since 2015, a human right and a social concern.^[1]

The current methods used to assess the maternal mortality ratio are very inaccurate and the numbers provided are not enough to recognize the major failures in diagnosis and treatment. In order to decrease the number of these errors the public health surveillance needs a

continuous interpretation of actualized data. Reliable data on maternal mortality implies an active vigilance, which requires the study of each new case. This would ideally involve an autopsy (clinical or forensic) as the circumstances demand. ^[3]

According to the DGS (Directorate-General of Health), in 2013, which are the last numbers in Portugal, maternal mortality is 8 deaths per 100000 live births. ^[5] This data is based fundamentally in clinic elements and in almost every case, the autopsy (clinical and forensic) is not included. ^[5]

The autopsy results are very important, because there are great discrepancies between the publish data and the real causes of death, that only can be confirm with the autopsy. This discrepancies became clear in 2008 in United Kingdom when it was performed a study with exhaustive evaluation, which included a series of autopsies. Unexpectedly, they realized, comparing with other countries, that indirect causes are more prevalent than direct causes. ^[6]

In 2009, a study in India proved that the discrepancy between clinical and autopsy diagnosis is more than 15%. ^[3] We believe that it is important to give more attention to the forensic study of maternal deaths in Portugal, in order to improve the published data and, therefore, defining health policies and be able to decrease the number of deaths.

The forensic assessment should include the clinical records, macroscopic and microscopic findings and toxicology, biochemistry and microbiological analyses. In order to do a complete autopsy study, it is important to keep in mind the specific and related pathologies, and some pathologies that can get worse with pregnancy. ^[3]

In maternal mortality it's known that there are four distinct categories, direct and indirect causes, coincidental and late deaths. ^[5]

This paper reviews the most recent published articles and the last statistical studies published by DGS and WHO, based on maternal mortality and the main direct and indirect death causes that can be neglected by doctors, leading to sudden death.

The aim is to create a review of the main pathologies that can affect pregnancy and to become a good tool for health care service, since almost every cause is preventable.

Therefore, we approach the definition, physiopathology, epidemiology and risk factors, clinic, diagnosis, and autopsy findings of each one. This work also aims to understand the role of the autopsy in sudden death during pregnancy and its possible contribution to the improvement of health care.

Materials and methods

Using article databases such as PubMed and books of obstetrics and legal medicine and organizations of the last ten years, we collected and integrated information about the etiology of maternal sudden death including the definition, physiopathology, epidemiology and risk factors, clinic, diagnosis, and autopsy findings. Maternal mortality can be divided into four categories, direct causes, indirect causes, coincidental and late deaths.

1) Maternal deaths directly related to pregnancy

Maternal deaths directly related to pregnancy include all deaths caused by obstetric problems. ^[7] The most frequent direct causes of maternal mortality are thrombosis and thromboembolism, pre-eclampsia and eclampsia, hemorrhagic disorders, amniotic fluid embolism and sepsis. ^[7]

1.1) Thrombosis and thromboembolism

Thromboembolism (VTE) is the combination of two pathologies: deep venous thrombosis (DVT), which is a condition caused by the formation of clots in the deep veins, usually in the

lower limbs (deep veins of the legs or in the pelvic veins, like the uterine veins); and pulmonary embolism, which is defined by the obstruction of lung arteries caused by clots.^[8]

In the antepartum period, the pathophysiology of VTE can be explained by: (a) the venous stasis caused by the progesterone-induced venodilatation, pelvic venous compression by gravid uterus and pulsatile compression of the left and right iliac veins, (b) hypercoagulability associated with increased thrombin production, and (c) fibrinolysis reduction due to the increased plasminogen activator inhibitor type 1 and 2 activity and decreased tissue plasminogen activator activity.^{[8] [9]}

In postpartum period, the vascular damage to the pelvic vessels that can occur after normal vaginal, assisted vaginal or cesarean section deliveries, and postpartum move restriction are the main alteration in the pathophysiology of VTE during the postpartum period. The hypercoagulability starts to decrease in the postpartum period and the risk gradually return to basal.^[9]

Embolism is one of the main causes of death in pregnant woman with an estimated risk of 0,8-4,7 per 100000 maternities.^[9] The highest risk of thrombosis starts with pregnancy and lasts until about two months after delivery.^[8] The main risk factors to VTE are: obese pregnancy, above 35 years old, sedentary life and caesarean delivery (rather than vaginal delivery) and coagulation mutations like the factor V Leiden.^{[7] [8] [10]}

The presence of symptoms like breathlessness, syncope, pleuritic pain, cough and hemoptysis, accompanied by the presence of signs like tachypnea, tachycardia and cyanosis should arise attention of the medicine assistant to the possibility of VTE.^[10]

The diagnosis of VTE is difficult to execute and normally only 10% have a final diagnosis. D-Dimers test, ultrasound, Doppler ultrasound, computed-tomography (CT), magnetic resonance (MR) or venography, are the most important support methods of the diagnosis.^[11] The American Thoracic society and the society of Thoracic Radiology have proposed an algorithm for the diagnosis of pulmonary embolism. This algorithm included CT pulmonary angiography, ventilation-perfusion scintigraphy, MR angiography and intravascular pulmonary angiography.^[11]

The review of the societies guidelines about thromboembolism prophylaxis (The American Congress of Obstetricians and Gynecologists, 2015; The American College of Chest Physicians, 2015 and The Royal College of Obstetricians and Gynecologists, 2015) shows that there is a significant difference between each others guidelines, but, even with an amount of studies conducted, they concluded that the thromboembolism risk during delivery has increased in the last years. They justify it with an inadequate risk assessment, inadequate thromboprophylaxis, failure to investigate symptoms, and failure to ensure multidisciplinary care.^{[12] [13]}

In cases of sudden death, if the autopsy is performed, it is necessary to check thrombus in the main pulmonary arteries to confirm the diagnostic of pulmonary thromboembolism. It is also important to confirm that the embolism was the cause of death and not a postmortem event.^[8] Then it is required the research of the thrombus origin which is made by dissecting the deep veins of the legs and the pelvic veins.^[8]

Hypertensive syndromes remain the leading causes of maternal morbidity and mortality. These syndromes have pathological consequences in the normal liver function.^[14]

The hypertension-related liver disease during pregnancy includes pre-eclampsia (PE), eclampsia (E), hemolysis, elevated liver enzymes, low platelets syndrome (HELLP) and acute fatty liver of pregnancy (AFLP).^[14]

1.2) Preeclampsia and eclampsia

Pre-eclampsia (PE) is an hypertensive syndrome characterized by hypertension ($>140/90\text{mmHg}$) plus proteinuria (upper urinary excretion to 0.3g of protein in a 24-hour urine sample) after the 20th week.^[8]

PE can be allocated into two categories, early and late PE. Although the etiology of this pathology is yet unknown, the main hypothesis in early PE cases is an alteration on the normal implantation of the placenta^[8] which includes the presence of an incomplete and imperfect invasion of the spiral arteries into the trophoblast during the implantation.^{[15] [16] [17]}

The later PE is related with an enlarged placental mass or surface. It can occur in cases of pregnant women with diabetes, nulliparous pregnant, anemia or in high altitudes.^[17]

The hypertension has major systemic repercussions, the most important of which is in the cardiovascular (increased in blood pressure and in cardiac output) and in hematological system (risk of hypovolemic shock, thrombocytopenia and disseminated intravascular coagulopathy (DIC)).^[15] Placental abruption, HELLP syndrome, thromboembolism and pulmonary edema, are some examples of the complications caused by hypertension progress.^[18]

Eclampsia (E) is an evolution of PE, with major neurologic changes like convulsions and coma, when other causes, such as epilepsy, are excluded.^[8]

Hypertensive syndromes (PE, E, HELLP syndrome) affect 2-8% of the pregnant women. Nulliparous pregnant, extreme age groups and family history have an higher risk of developing PE. Higher risks are also associated with systemic lupus, diabetes and chronic hypertension.^[6]

Typical signs and symptoms of hypertensive syndromes are: headache, epigastric pain, visual alterations, oliguria or anuria, hepatocellular dysfunction and hyperreflexia.^[16]

During pregnancy, the diagnosis can be done by periodically measuring the blood pressure and the proteinuria (screened with 24-hours urine test).^[8] The symptomatology, the presence of high blood pressure ($140/90\text{ mmHg}$)^[11] and upper urinary excretion to 0.3g of protein in a 24-hour urine sample, make the diagnosis of pre-eclampsia.^[8] Nowadays, some studies about the substitution of 24-hours urine collection for the 12-hours test show an advantage of convenience and improved clinical efficiency (of the later). Furthermore, there are investigations in order to implement some biomarkers in the first and second semesters to prevent this syndrome from de beginning. For the first semester it is proposed Placental Growth Factor (PIGF) and in the second semester the Human Chorionic Gonadotrophin ($[\text{hCG}]>2.0\text{ MoM}$) and Associated Plasma Protein-A ($[\text{PAPP-A}]<5\text{th centile}$). These biomarkers seem to have more accuracy to predict pre-eclampsia. The importance of these new biomarkers is the need for early start of tracing PE. Nowadays this prevention only starts in the mid to the end of the second trimester and the dysfunction of the placenta may already be establish.^{[17] [19]}

The large discordance between the cutoff of the different studies is the greatest obstacle to the progress in prevention and early diagnosis of hypertensive syndromes.^[17]

PE kills 3 women per day, this remains as one of the most frequent causes of death in pregnant women. In the case of sudden death the main findings at autopsy are a soft liver parenchyma with associated hematoma or a large haemoperitoneum caused by the liver capsule's rupture especially the right lobe (liver rupture).^[7] The microscopic section shows edematous changes in fibrinoid material, and recent blood extravasation.^[8] A third of the cadavers present brain alterations, with subcortical edema, vascular lesion consistent with fibrinoid necrosis of arterial walls, perivascular infarction and hemorrhage within the white matter, basal ganglia and pons or even a ventricular rupture can be present.^[11]

1.3) HELLP syndrome

The HELLP syndrome is characterized by hemolysis (consequent of a microangiopathic hemolytic anemia (MAHA)), elevated liver enzymes and decreased platelet concentration ($<100.10^9/L$).^[20]

This syndrome is classified as a severe variant of the PE and it occurs in about 5 to 10% of pregnant women with severe PE^[8] with a frequency of 0,2-0,6% during pregnancy.^[14] The peak of incidence is between 27 and 37 weeks and until 48 hours after delivery.^[20] Contrary to PE, the risk factors are multiparous pregnant women, caucasian and older than 35 years old women.^[16] In a posterior pregnancy the risk of recurrence is 20%.^[20]

The main symptom of HELLP syndrome is an upper right quadrant pain or epigastric pain, that worsens overnight and can be associated with nausea and vomiting. These symptoms usually progress continuously but the intensity can change spontaneously.^[20]

The diagnosis include laboratory analysis with platelets count under $<100.10^9/L$ and an increased concentration of liver enzymes (AST, ALAT). The last analysis, often reflects the presence of hemolytic processes. Although the increased levels of α -GST and GST-a1 are the most sensitive factors to appoint acute liver injury, they cannot be used in diagnosis yet.^[20]

HELLP syndrome have a maternal mortality of 3,4-24,2 %.^[21] The autopsy findings are similar to preeclampsia.^[8]

1.4) Acute fatty liver of pregnancy

Although still poorly elucidated, the main hypothesis to explain this condition is the deficiency of the long-chain 3-hydroxyl-CoA dehydrogenase.^[8] This alteration interferes in the fatty acid degradation through beta oxidation in hepatic mitochondria.^[14]

Acute fatty liver of pregnancy is a potentially fatal condition but it is very rare with an incidence of 1 per 10000 pregnant women.^[3, 8, 14, 16] It occurs most frequently in the third trimester and the differential diagnosis with other hypertensive syndromes can be difficult.^[16]

The first symptoms related with this pathology are headache, nausea, vomiting, abdominal or focalized right upper quadrant pain, itching and confusion, which lead the pregnant women to the hospital.^[11]

The diagnosis is difficult and normally the first diagnostic hypothesis are PE or hepatitis. Although biopsy is rarely done due to the large risk to the pregnancy, it is the only way to confirm the diagnosis. Imagiological exams like ultrasound and magnetic resonance have low sensibility. That's why the presumed diagnosis is done by the clinical and laboratory findings (elevation in the transaminase, direct bilirubin, and uric acid levels, anemia normocytic normochromic and leukocytosis)^[16]

The major complications and causes of death in pregnant women, with acute fatty liver, are renal failure, DIC and sepsis.^[16] At autopsy, it is possible to check a fatty liver.^[8] The diagnosis is microscopically confirmed with the presence of microvascular centrilobular steatosis, cholestasis, cytoplasmic ballooning, inflammation and necrosis of hepatocytes.^[8]

Obstetrical hemorrhage continues along with infections and hypertension to constitute the famous triad responsible for the majority of cases of death in pregnant women. The decreasing maternal mortality rate related with hemorrhagic causes has as been achieved in recent years and that represents a great global maternal mortalities decrease.

1.5) Hemorrhage

1.5.1) Hemorrhagic stroke

Hemorrhagic stroke (HS) incidence has been increasing lately. It is subdivided in subarachnoid hemorrhage (SAH) and intracerebral hemorrhage (ICH). The SAH results, more frequently (approximately 80%), from a cerebrovascular alteration or a saccular aneurysm rupture.^[11, 22] In a lower percentage, SAH can be a consequence of a coagulopathy, angiopathy, venous thrombosis, infection, drug abuse, cancer or trauma.^[11] The ICH is caused by a rupture of a small vessel previously damage by chronic hypertension.^[11] Although a stroke is uncommon during pregnancy or the postpartum period, a venous infarction thrombosis of dural sinuses can cause it. It has a multifactorial etiology including hypercoagulable state typical of pregnancy, protein S deficiency and dehydration.^[8]

HS has a prevalence of 6 per 100000 deliveries.^[22] The advanced age (above 35 years old), arterial hypertension, pregnant woman medicated with anticoagulants, antiplatelet and fibrinolytic drugs, alcoholism and the presence of arteriovenous malformations are the major risk factors recognized.^[8] During the postpartum period the risk of SAH increase.^[8]

The major symptoms and signs are headache, vomiting and local neurologic deficits.^[8] The diagnosis of this condition implies a large index of suspicion by the clinician. A lot of pregnant women enters the emergency service with few classic risk factors of HS or only with a slight increase of the blood pressure.^[22] Imagiologic exams after suspicious clinical findings are the best way to determine the diagnosis of SAH.^[8]

The mortality percentage of SAH and ICH are 10% and 20% respectively.^[22] At autopsy it is possible to evaluate the presence of hematoma. Microscopic section may present vascular malformations or other vascular disease, infection findings, cancer and other possible etiologies.^[8] The autopsy can confirm the presence of thrombosis in the dural sinuses.^[8]

1.5.2) Haemoperitoneum

Facing haemoperitoneum the clinician should consider a uterine rupture, ruptured subcapsular hematoma, aneurysm rupture of the splenic artery, ectopic pregnancy, placenta previa, placenta increta and abruption placentae.^[8]

a) Ectopic pregnancy

Ectopic pregnancy is the implantation of the blastocyst outside the endometrial lining of the uterine cavity. This occurs in consequence of the anatomic or functional obstruction, impair tubular motility, ciliary dysfunction and molecular chemotactic factors that stimulates and protects tubular implantation.^[23]

Although this is a rare cause of maternal death, is still the primordial cause in the first trimester due to hemorrhage and infection, with a prevalence of 20 per 1000 pregnancies.^[16] The most frequent location is in the fallopian tubes (98%).^{[8] [16]}

Genital and pelvic infections are risk factors of ectopic pregnancy, as the use of medically assisted reproduction techniques, tobacco, scar of previous surgery and age above 35 years old.^[16]

A pregnant woman may have an acute or subacute symptomatology. The most common presentation is a woman with a menstrual delay, positive pregnant test, abdominal-pelvic pain, metrorrhagia and hypovolemic shock.^[8] More than 50% of the pregnant women with ectopic pregnant have at least a risk factor.^[23]

The diagnosis of this pathology includes reunion of clinical findings, transvaginal ultrasound and B-HCG ($>1000\text{IU/L}$) dosing.^[23] If the diagnosis is not clear, an exploratory laparoscopy can be done.^[8]

A tubal ectopic pregnancy location can result in tubal abortion, tubal rupture or spontaneous resolution.^[15]

At autopsy it's possible to observe the ectopic localization and the haemoperitoneum in case of rupture.^[8]

b) Uterine rupture

The uterine rupture (UR) is a disruption of the uterine wall that can be total or partial.^[8] UR can be traumatic or non-traumatic.

UR has a prevalence of 1 per 100 deliveries to 1 per 1850 deliveries.^[8] In non-traumatic cases, the risk factors are previous caesarean section, increased gestational age, multiparity and the use of uterotonic drugs (like oxytocin and prostaglandins). Uterine rupture can occur during an obstetric intervention as mid-forceps delivery and prolonged labor with cephalopelvic disproportion.^[8] The traumatic uterine rupture is due, in most of the cases, to road accidents.^[8]

In the beginning of pregnancy, women may have an acute abdomen frame with intense pain and signs of internal bleeding. With the advance of pregnancy, the patient may be asymptomatic or only presenting a slight abdominal pain or vaginal bleeding. During labor, the rupture is felt with a strong localized pain in the abdomen.^[16]

The diagnosis is based on clinical history and physical examination. In case of doubt, it is possible to resort to an ultrasound to confirm the diagnosis.^[16]

In the autopsy it's possible to observe the haemoperitoneum and the rupture of the uterine wall.^[8]

c) Placenta implantations

The placental previa (PP) is defined as a low implantation of the placenta that obstruct totally or partially the internal os. Low-lying placenta is defined as a low implantation localized at least two centimeters of the internal os.^[11]

PP is an important cause of bleeding during the 3th trimester of pregnancy and delivery. This pathology has a prevalence of 3 to 5 per 1000 deliveries.^{[24] [20]} The major risk factors are multifetal gestation, age above 35 years old, previous cesarean section, cigarette smoking and an increase in perinatal screening of maternal serum Alfa- fetoprotein.^[11]

The most typical and frequent sign is a moderate painless vaginal bleeding.^[11]

The diagnosis is made by ultrasound or MR, allowing verification of the placenta position.^[16]

In the autopsy it's possible to confirm the wrong position of the placenta.^[8]

The excessive invasion of the myometrium by the placenta is called accreta. Different degrees of invasion have different classifications (accrete, increte, precreta). The accrete placenta is the main cause of intractable postpartum hemorrhage and emergency hysterectomy intrapartum.^[24]

The defect of the decidua with an imperfect fibrinoid or Nitabuch layer and the alteration of cytotrophoblasts, that can control the decidua invasion through the control of angiogenesis and growth factors, represent the major alteration in this pathology.^[11]

The incidence of accretism is 1 per 533 deliveries. Throughout time it was discovered a correlation between increte placenta and placenta previa, previous uterine curettage, previous cesarean section, multiparity and advanced maternal age (above 35 years-old).^[24]

Typically, pregnant women resort to emergency service with vaginal hemorrhage that may

appear only in the third trimester.^[11]

Although the diagnosis can be made by Doppler ultrasonography and MR imaging, when a pregnant woman is asymptomatic (absence of vaginal bleeding) these tests are less sensitive.^[24]

The accrete placenta represents 8% of the deaths related with hemorrhage. In autopsy, the anatomopathologic diagnosis of accretism is made by histological study with the presence of trophoblastic islands in myometrium.^[8]

The abrupt placenta (AP) is the separation, total or partial, of the placenta from the local of implantation. AP begins with hemorrhage to the decidua that then split, leaving a thin layer adherent to the myometrium. The etiology can be an impaired trophoblastic invasion with subsequent atherosis. Other possibility can be inflammation or infection of the myometrium.^[11]

The prevalence is 4 to 35 per 1000 deliveries.^[25] AP can be a cause of dangerous bleeding in addition to the risk of intrauterine death.^[20, 24] This injury can be caused by a traumatic event or it may have a non-traumatic cause like maternal hypertension, cigarette smoking and cocaine abuse.^[8]

The main symptoms and signs are abdominal pain and abdominal tenderness. The shock can manifest within few minutes or hours after the injury.^[8, 24]

The diagnosis is based on clinical history and physical examination and supported by some imaging tests such as ultrasound and MR.^[8, 25] In the initial phase pregnant women can be asymptomatic, if there isn't more abruption of the placenta, the diagnosis can only be made after delivery when assessing the placenta, when the clinician verifies an area few centimeters wide with a dark surface and with blood clots.^[11]

At autopsy, the placenta must be verified in order to find signals of abruption like dark surface and the presence of blood clots. The presence of cocaine in the blood and the presence of signs of trauma should be analyzed too.^[8]

d) Ruptured aneurysm of the splenic artery

The ruptured aneurysm of the splenic artery is more common in the third trimester of pregnancy and this pathology has an high mortality rate. The main causes are unknown, but the main hypotheses are hemodynamic and hormonal (increase in the estrogens and progesterone concentrations) alteration during pregnancy, uterus compression and the increase in the arteries flow.^[26]

The increase in estrogens and progesterone concentration can have as consequence the disruption of the internal elastic lamina, fragmentation of the elastic fibers, degeneration of smooth muscle fibers, and failure of elastin formation. The relaxation can be increased too and that may affect the elasticity of the splenic artery wall.^[26]

The majority of this aneurysms are saccular, located 2-5 cm from the splenic hilum.^[8] The prevalence is between 0,01 and 10,4 %. The multiparity, portal hypertension, atherosclerosis and medial fibrodisplasy increase the risk of this pathology.^[26]

Pregnant females who have this condition may be asymptomatic (when the aneurysm has less than 2 cm in the major diameter or they can present an epigastric pain with or without signs of shock.^[27] The diagnosis has improved recently due to the evolution and use of imaging studies as angiography, the MR and CT.^[27]

The mortality of the ruptured aneurysm of the splenic artery is around 75%.^[28] At the autopsy, it can be very difficult to identify the aneurysm. It is important to carefully dissect the full length of the splenic artery.^[8]

Amniotic fluid embolism (AFE) is a rare cause of sudden death in pregnant women, but, it is considered by many, as a key factor.^[29] Mortality has decreased by raising awareness as a condition that can become catastrophic.^[29]

1.6) Amniotic fluid embolism

Despite the lack of information, it is thought that the main cause of AFE is related to immunology, where an idiosyncratic reaction is triggered by the access of amniotic fluid to the maternal circulation. This access implies failure in the physical barriers in utero-placental unit (e.g. Endocervical veins).^[29] Recurrence risk is very low.^[29] The coagulopathy occurs as a result of both procoagulant and anticoagulant factors and it's likely to be multifactorial.^[29]

The incidence, 7 to 8 per 100000 births, is still somewhat vague due to the different types of methodology used in studies and the low number of studies.^[29]^[8]^[30]

The major risk factors are the changes in the barriers (e.g., placental abruption) or increased intrauterine pressure (e.g., uterine contractions), maternal age above 35 years old, multiple pregnancy, cesarean section, assisted delivery or eclampsia.^[29]

Clinical signs related with AFE are hypotension, fetal distress, pulmonary edema or ARDS, cardiopulmonary arrest, cyanosis and coagulopathy. However, the clinical presentation usually has a great diversity of signs and symptoms.^[29]

Diagnosis is based on four inclusion criteria (1) the acute hypotension or cardiac arrest; (2) acute hypoxia; (3) severe coagulopathy with clinical bleeding in the absence of other explanations; (4) an event during delivery, cesarean delivery, surgery to the uterus or within a period of postpartum and defined by no other explanation for the results.^[29] Imaging study can give some contribution to the diagnosis, but there is no specific test.^[29]

The diagnostic can be achieved by demonstrating the presence of squamous cells of the fetus in maternal blood, through the introduction of a catheter into the pulmonary artery.^[8]

Clinical suspicion, traditional laboratory data, or intravascular cellular debris (demonstrated only in 50% of patients) are insufficient to make a definitive diagnosis of AFE. Nowadays, biomarkers capable to confirm the diagnosis are inexistent. Modern biomarkers may help differentiate AFE from other conditions.^[30]

The highest levels of STN (Sialyl Tn), an fetal antigen present in meconium and amniotic fluid, which is an high sensitivity test used, were found in AFE non-survivors and can be the key for predicting the severity and mortality of AFE.^[29]

The main consequences of this pathology are heart failure and DIC.^[31] In cases of sudden death it's hard to find the failure of the barrier.^[8] The autopsy diagnosis is made by histological studies with the detection of fetal products in the maternal pulmonary arteries, like squamous epithelial cell, meconium, lanugo and mucins (origin in the intestinal tract of the fetus). The difference between fetal cells and maternal squamous cell shedding is made by the presence of nuclei in the maternal peeling endothelial cells, which normally does not appear in fetal squamous cells.^[8] We can still do immunohistochemical staining with cytokeratin that mark the squamous cells but not the endothelial ones. Mucins are usually present but it is hard to detect because they rarely stain with routine dyes. However it's possible to stain with "Alcian blue" or mucins staining. It is sometimes possible to find products of amniotic fluid in the uterine and cervical veins.^[8]

Sepsis remains a leading cause of maternal death, in contrast with what was expected, the maternal sepsis has increased even in developed countries.^[32]

1.7) Sepsis

Nowadays the main scientific societies have different definitions for the classification of sepsis. This fact makes the compression of this pathology even harder. ^[32]

WHO, in 1992, defined puerperal sepsis as an infection of the genital tract occurring at any time between the rupture of membranes or labour and the 42 days in the postpartum period. Two or more of the following symptoms are present: pelvic pain, fever, abnormal vaginal discharge and delay in the reduction of the size of the uterus. ^[33]

ICD-10 (International Classification of Diseases) defines puerperal sepsis as an higher temperature of 38°C continued for 24h or recurrent during the period between the end of the first day and the end of the tenth day after birth. ^[32]

In 2002, a surviving sepsis campaign defined systemic inflammatory response syndrome (SIRS) as a widespread inflammatory process caused by an infection, trauma, thermic lesion or sterile inflammatory process. It requires the presence of two of the criteria: (a) temperature <36°C or >38°C, (b) frequency cardiac >90 bpm, (c) respiratory frequency >20cpm and (d) leukocytes <4 x 10¹⁰ cells per liter or > 12x 10⁹ cells per liter. ^[7]

The surviving sepsis campaign defined sepsis as a SRIS associated with an infection. ^[32]

Severe sepsis presents hypotension and hypoperfusion, which responds to fluid resuscitation, and organ dysfunction. And finally septic shock is the worst state of sepsis without fluid resuscitation response ^[7].

Sepsis may have an obstetric or non-obstetric etiology. Obstetric causes include septic abortion, chorioamnionitis, perineal infection, endometritis, surgical wound infection and puerperal mastitis. The non-obstetric causes included urinary infection, respiratory infection, appendicitis, pancreatitis, hepatitis, malaria and infection by the human immunodeficiency virus (HIV), among others. ^[32]

Between all this causes, pyelonephritis, chorioamnionitis and endometritis are the most important causes of septic shock in pregnancy and postpartum period. ^[32]

An american study, conducted between 1998 and 2008, showed a sepsis incidence of 1 per 17246 births. The most frequent risk factors depend on the infection agent, host factors and pregnancy and delivery factors. ^[32]

Anemia, obesity, previous caesarean section, premature rupture of membranes, diabetes mellitus or glucose intolerance, immunosuppression, tobacco, maternal age above 35 years old, chronic diseases and low socio-economic stratum are included in the host factors. ^[32]

The main agents related with maternal death are streptococcus group A beta-hemolytic and *Escherichia coli*. The first agent exists in 5-30% of the population as an asymptomatic way. ^[7]

The main clinical findings are fever or hypothermia, tachycardia, tachypnea, hypoxia, hypotension, oliguria and state of consciousness. ^[32] The diagnostic is done by the clinical findings, a blood culture and C-reactive protein. The signals and symptoms can be subtle due to the physiological adaptation of the pregnancy. ^[32]

Blood cultures should be made in the source of infection such as vaginal, respiratory secretions or cerebrospinal fluid before the introduction of antibiotics. Imaging scan like chest x-ray, ultrasound and computer tomography may help the diagnosis. ^[7]

Sepsis in pregnant women can result in increased preterm birth rate, hypoxia, acidosis, stillbirth and fetal infections. ^[34]

Puerperal sepsis has worldwide mortality of 15%. The recent increase in maternal deaths in developed countries can be justified with (a) the advanced maternal age, obesity, cardiovascular disease, chronic renal diseases, erythematous systemic lupus and smoking; (b) household childbirth; (c) increased number of infections by Streptococcus beta Group A beta-hemolytic and *Escherichia coli*; (d) the use of assisted reproductive techniques, invasive prenatal diagnostic test, and caesarean delivery ^[32]. An antibiotic prophylaxis can reduce the incidence of sepsis and, consequently, the incidence of maternal deaths. ^[32]

Sepsis rarely is the primary diagnosis. Normally the objective of the autopsy is to demonstrate a concomitant disease, which may have been neglected. The best interval to do the autopsy is in the first 15 hours after death and the acquisition of some clinical information is fundamental before starting the post-mortem exam.^[35]

Sampling may change from situation to situation but the more extensive and varied it is, the possibility to do the diagnosis and determine the starting point of the infection increases.^[35]

Examples include sampling of blood, organs product aspiration, cerebrospinal fluid, urine and feces. The culture of these products is still very effective, however, nowadays other exams including DNA technique like PCR supported by immunohistochemistry can facilitate the diagnosis.^[35]

2) Maternal deaths indirectly related to pregnancy

Maternal deaths indirectly related to pregnancy include all deaths caused by the intensification of a disease that already existed prior to pregnancy.^[7]

The cardiac diseases are causing more deaths in recent years. It includes some examples like peripartum cardiomyopathy, aortic dissection, arrhythmogenic right ventricular cardiomyopathy among others. Central Nervous System (CNS) disorders most frequent are intracerebral hemorrhage, cerebral thrombosis and epilepsy.^[7]

Also included respiratory diseases (e.g. asthma), endocrine, metabolic and immunologic diseases (e.g. diabetes), gastrointestinal system diseases (e.g. pancreatitis), blood disorders (e.g. thrombotic microangiopathy), circulatory system diseases, kidney and tumors (e.g. pheochromocytoma).^[7]

The HIV or bacterial infections prior to pregnancy are also indirect causes.^[7]

2.1 Cardiac disease

The cardiac disease in pregnant women may be one of the main indirect causes of sudden death.^[7] This pathology has a prevalence of 2,31 per 100000 pregnancies.^[7] The main risk factors are obesity, tobacco and hypertension.^[7]

The etiology can be congenital (pulmonary hypertension and cardiac congenital diseases) or acquired (ischemic and non-ischemic).^[7]

The acquired non-ischemic conditions include the aortic dissection, peripartum cardiomyopathy, myocarditis, arrhythmogenic right ventricular cardiomyopathy, among others.^{[8][7]}

2.1.1) Peripartum cardiomyopathy

Peripartum cardiomyopathy (PC) etiology, like most of the disorders during pregnancy, is still under research.^[8] The main hypothesis for PC is a fetal autoimmunity and microchimerism, myocarditis and the dietary excess of salt or deficiency of selenium. PC is marked by the development of systolic heart failure.^[36]

PC can have an hereditary and genetic component.^[36] The most recent studies propose a truncating variants as a strong genetic predisposition.^[36]

This cardiomyopathy it's more frequent in the last month of pregnancy until 6 months after birth, with the peak of incidence in the second month after delivery.^[8]

The incidence of this pathology is 1 per 1000 or 4000 births.^[8] ^[36] PC is more frequent in black pregnant women, age above 35 years old, multiparous, preeclampsia and hypertension.^[7]

Pregnant women with this condition usually present dyspnea, edema, orthopnea, tachypnea

and tachycardia.^[7] The diagnosis is made with a chest x-ray and an echocardiogram. They should be accompanied by a specialist and possibly included in the regional transplant center.^[7] In US, PC represents 4% of all cardiac transplants.^[36]

PC has a mortality of 5 to 10%, and it is a rare cause of maternal sudden death.^[36] The most frequent mechanisms of death are attributed to arrhythmia or thromboembolism.^[8] At autopsy there are nonspecific findings as enlarge heart, dilated cardiomyopathy, irregular myofibrils and fibrosis, and variable myocardium T cell infiltration.^{[7][7] [8]}

2.1.2) Arrhythmogenic right ventricular cardiomyopathy

Arrhythmogenic right ventricular cardiomyopathy (ARVC) is an inherited disease in which 50% of the cases with autosomal dominant transmission. ARVC is characterized by fibro-fatty changes of the right ventricle, ventricular arrhythmia, and sudden death.^[37]

Although this ARVC is normally considered a pathology of desmosomes, in fact only 50% of the cases have mutations in the genes of this component. Recent studies in mice showed changes in signaling via Rho-kinase, which is implicated in the regulation of intracellular transport and cytoskeletal filament organization, that supports the desmosomal complex proteins in cell-cell junctions in the myocardium. Mice that had Rho-kinase inhibition exhibited spontaneously developments of cardiac dilatation and dysfunction, fibro-fatty change and ventricular arrhythmia that resulted in premature death of the mice. This pattern corresponds with ARVC in humans.^[37]

Ventricular tachycardia is provoked by hemodynamic changes of pregnancy, as the increased in the cardiac output or anemia when associated with prior cardiac pathology.^{[38][39]} This pathology has an incidence of 1:5000 births.^[38]

This disease has a ranging spectrum of presentation, from asymptomatic to sudden death. Usually pregnant woman presents with palpitations and syncope. The diagnosis is made by electrocardiogram, classically with a left bundle-branch block that can be accompanied by inversion of the T wave from V1 to V3 and premature ventricular complexes.^[38] Nowadays the availability of genetics tests increases the early recognition of ARVD.^[38]

In the autopsy it's possible to evaluate the fibrous-adipocyte tissue replacement myocardium, although it is not pathognomonic. Normally there is attenuation of myocytes, transversal stretch marks and areas with less myofibrils, this alteration gives the appearance of "bubbly".^[40]

2.1.3) Aortic dissection

Acute aortic dissection (AAD) is a rare phenomenon during pregnancy despite a well-established risk relationship.^[41] AAD is characterized by a rupture of the intimal layer of the aorta with blood extravasation, forming a subintimal hematoma that produces a false double-channel effect.^[41] It has an incidence of 1,39 cases per 100,000 person/year among pregnant women under 40, compared to 0.4 cases per 100,000 person/year in pregnant women aged between 15 and 45.^{[41] [7]} As previously mentioned, pregnancy is a risk factor, which is increased with gestational hypertension, preeclampsia, Marfan and Ehlers-Danlos syndrome.

Pregnant women with sudden and persistent thoracic and interscapular pain should lead to an AAD diagnosis that can be done by Trans-esophageal echocardiography, a CT scan or an MRI.^{[7][41]}

AAD can complicate with aortic valve insufficiency, avulsion of the carotid arteries resulting in myocardial infarction or stroke.^[41] This pathology represents 3-14% of maternal cardiac death.^[41]

In autopsy it is possible to observe hemothorax and a rupture of the aortic or coronary artery.^[8] Microscopically it's possible to confirm the recent artery dissection with the underlying aortic medial degeneration. Because Marfan and Ehlers-Danlos syndromes are both great enhancers of aortic dissection as in global population as in pregnant women, genetic testing for these syndromes should be done.^{[41][7]}

2.1.4) Cardiac ischemia and myocardial infarction

Nowadays, myocardial infarction is one of the most frequent causes of pregnant cardiac death (20%). The etiology can be variable like atherosclerosis, coronary dissection, or even an autoimmune disease.^[7]

In more than 25% of the cases the main etiology is coronary dissection, most frequent in the left descendent anterior artery, in the anterior wall (78%).^[42] The coronary dissection is more frequent in the postpartum period and it is possibly a result of the hemodynamic and hormonal changes during pregnancy. In young pregnant women the most frequent etiology is embolic and normally with classic risk factors but with the presence of structural cardiac diseases.^[42]

Coronary spasm and dissection can be consequent of drug abuse (e.g. cocaine).^[42]

The prevalence of cardiac ischemia (CI) is 6.2 per 100.000 pregnancies.^[43] CI is more frequent during the third trimester and six weeks following the deliver.^[43] The risks factors are obesity, tobacco, age above 35 years old, more than 3 births, diabetes, hypertension, thrombophilia and family story.^{[7][43]} Pregnancy increases three or four times the risk of a CI.^[43] In pregnant women, and because the etiology is rarely atherosclerosis, the symptoms can be very different. The most frequent ones are abdominal or epigastric pain associated by vomiting and dizziness.^[7]

The diagnosis is made by a serial of parameters that include the symptomatology, electrocardiogram (ECG) and the study of the cardiac enzymes (like troponin). A normal ECG does not exclude the diagnosis. In order to overtake this problem, women should do various ECG. The study of coronary arteries is also necessary trough a coronary angiogram with angioplasty with or without stenting.^[7]

At autopsy it is possible to evaluate the typical macroscopic and histological changes of a myocardial infarction and also explore the etiology. In the majority of the cases, it's possible to find atheroma plaques in the coronary arteries or a dissection of the same arteries.^[7]

2.2) CNS (central nervous system)

2.2.1) Epilepsy

Sudden death in epilepsy (SUDEP) is defined, in pregnant population as in general population, as: *"death, unexpected, witnessed or unwitnessed, non-traumatic and without the patient drowning due to epilepsy, with or without evidence of a seizure and excluding the documented epilepticus status, where autopsy did not reveal a toxicological or anatomical cause of death."*^[44]

The physiologic alteration during the pregnancy, such as the increase in the progesterone and estrogen and the alteration in the absorption, distribution, metabolism and elimination of the antiepileptic drugs, is the main hypothesis for the negative impact of epilepsy.^[45]

In many countries such as the UK, studies have concluded that the mortality rate increases 10 times during pregnancy in women with epilepsy when compared to women without the disease.^[44] The prevalence of epilepsy in pregnancy is 0.61 to 1 per 100,000 births.^[7] The major risk factors of SUDEP in a pregnant woman are chronic epilepsy with poorly controlled seizures, hyperventilation, sleep privation, stress, physical pain and lack of compliance with

medication.^[46]

Sometimes it is difficult to diagnose epilepsy during pregnancy due to the proximity of clinical signs typical of pregnancy. Headaches, changes in mood, dizziness, confusion and memory loss are some of the most common symptoms.^[46] The diagnosis of epilepsy is mainly clinical but there are auxiliary diagnostic tests such as EEG, CT and MR that can help.^[47]

Guidelines recommend to prevent SUDEP *(a) choose the optimal treatment before conception, (b) prefer monotherapeutic approach if possible, (c) opt for the most efficient AED according to the type and frequency of seizures, (d) begin with the lowest effective dose, and (e) not forget folic acid supplementation.*^[45]

In autopsy of SUDEP cases it's possible to find alterations in several organs including (a) brain (edema and hypoxia signs), (b) heart (fibrosis driver system), (c) lungs (increased weight and pulmonary edema with alveolar hemorrhage and exudate) and finally the (d) liver (increased venous congestion and weight which are cardiac failure signals).^[48]

2.3) Blood flow

2.3.1) Thrombotic microangiopathy

Thrombotic microangiopathy include primary disease like thrombotic thrombocytopenic purpura (TTP) and hemolytic uremic syndrome.^[8]

TTP characterized by the presence of microthrombi rich in vWF and the presence of ADAMTS13 inhibitors in the blood.^[49] Normally, it arises before of 24 weeks.

The prevalence of this pathology is 1 per 25000 to 1 per 100000 pregnancies.^[50]

Most frequent symptoms and signs are neurological (lethargy, seizures, mental confusion), fever, kidney dysfunction, severe thrombocytopenia (gingival, gastrointestinal or genital bleedings, bruises, petechial), abdominal pain, diarrhea, anemia, intravascular hemolysis, visual disturbance and cardiac dysfunction by thrombotic events in the microcirculation.^[16]

In the case of pregnant women who have already TTP prior the pregnancy it's normal the worsening of the situation. Unfortunately, in many cases it is necessary to terminate pregnancy before fetal viability.^[16]

The absence of pathognomonic laboratory data hampers the diagnosis. Identification of ADAMTS-13 deficiency is helpful to allow specific care of this potentially fatal disease. When the diagnosis is done in postpartum period, the maternal prognosis is reserved.^[16] Lactic dehydrogenase (LDH) is always very high due to hemolysis.^[16]

TTP is a disease with a mortality rate higher than 60% when untreated and although it is not a specific disease of pregnancy, this pathology can be triggered by it.^[49] In cases of sudden death, at autopsy the conditions that cause the formation of widespread microthrombi are called thrombotic microangiopathy of pregnancy.^[8]

2.4) Tumors

2.4.1) Pheochromocytoma

Pheochromocytoma is a chromaffin tissue tumor located, in 80-85% of cases, in the adrenal glands and, in 15-20% of the cases, in extra-adrenal sites (for example Zuckerkandl body) characterized by increased release of catecholamines.^{[51] [52]} This tumor typically known as "10% tumor" because it is 10% bilateral, 10% multiple, 10% hereditary and 10%

malignant.^[51]

It is extremely rare in pregnant women, with an incidence of 1 per 50,000 pregnancies.^[51]

Clinically the main symptoms are headache, hypertension, hyperhidrosis, and heart palpitations.^{[51] [52]}

In pregnant women, despite the absence of screening tests, it is recommended to carry out biochemical tests, measurements in blood and urine catecholamine metabolites levels (metanephrine and Normetanephrine) when in presence of a sustained or paroxysmal hypertension, clinical signs of increased secretion of catecholamines or when they have a family history of pheochromocytoma.^{[51] [52]}

When biochemical tests have positive results, the second phase of diagnosis is achieved by undertaking imaging tests to locate the tumor.^[51]

The risk of sudden death in pregnant women with pheochromocytoma is due to paroxysmal hypertension and hypotension that may be precipitated by vaginal delivery, the fetal movements, abdominal palpation, uterine contraction, bleeding or general anesthesia.^[52] Vaginal delivery has been contraindicated in the past but is acceptable nowadays if the patient is stable.^[52]

The mechanism of death normally is necrosis of the tumor produced by the compression of the gravid uterus that leads to a cardiopulmonary failure by the extensive liberation of catecholamines.^[51]

At autopsy it's possible to find a mass in the adrenal gland that it is formed in the medulla layer. [51, 53] Macroscopic analysis demonstrate a necrotic and hemorrhagic mass. In microscopic analysis it's possible to confirm the endocrine origin with the presence of polygonal cell with low number of granules and acidophilic cytoplasm, decentered nucleus fusiform and round elements that suggest anisocytoses, as well the disposition of the cell over rows conjunctive and vascular tissue with an anarchic orientation.^[51]

Discussion and conclusions

Sudden death in pregnancy has a major impact in the society, but actually in Portugal, as in many other countries, it has been poorly explored.

In 2007, DGS published a report about the main causes of maternal mortality. Although these datas have more than 8 years, these were the last numbers presented in Portugal.^[54] This publication includes maternal mortality (MM) (figure 1), ratio of maternal mortality (RMM) (Table 1), the maternal death by age groups (Table 2) and the main causes of maternal mortality (Table 3).^[54]

The most frequent causes are cardiac disease such as, pulmonary hypertension, cardiac congenital disease, aortic dissection, peripartum cardiomyopathy, myocarditis, right ventricular arrhythmogenic cardiomyopathy, and ischemic disease.^[7, 54] A percentage of 21,7% of the maternal death had unknown cause.^[54]

In 2015, DGS publish a report with the most important health index, whose maternal mortality was included. This report show the datas of 2013, with a maternal mortality of 8 deaths per 100000 live births.^[5]

The last data publish by the WHO, in 2015, indicated that Maternal Mortalities are 12 and 239 per 100000 live births in developed and developing countries respectivel.^[2] With this data we can concluded that Portugal is below of the mean of the developed countries, even so, the importance of the reduction of this numbers is crucial.

Nowadays knowledge, allows the prevention of the majority of deaths, but failures in diverse points turn it not a reality yet. Failure to detect signs or symptoms and the late feature of the

pregnant women to medical cares are actually the two most important causes of maternal death.^[55]^[56] Although it is important to review the services provided to every pregnant, it is also important understand the various barriers that stand between pregnant woman and these services. The difficulty in transport to medical facilities, low educational level and economic problems are some of the obstacles for the preventive measures come evenly to all pregnant women.^[57]

Since all pregnant women, and especially those who have high risk from the beginning, are at risk of suffering complications that can lead to her death, it is important to implement measures that allow uniform access to maternal health care from the start and more intense in the third trimester as we already been mentioned, is the time when the risk of death increases.^[55]

The production of more recent reports that specifically show the current numbers of each cause of maternal death and the number of case that the etiology remained unknown is actually very important to understand the reality of our country in order to actualized our guidelines in the routine of the pregnancy vigilance and to create better access to the health care for the pregnant woman. So, to complete this reports it's very important that each case is study individually and, in the end, all cases have a definitive diagnosis of the cause of death. The diagnosis can be made by the clinical with de clinic symptoms, sings and auxiliary test. If the clinician has doubts about the pathology, it is possible to accomplish a clinic autopsy. On the other hand, if there are some clinical doubts it's necessary to perform a medical legal autopsy. The pos-mortem study should be standardized in order to avoid misclassification and loss of information.^[7] A study based on anatomo-pathological findings is a good possibility to better understand the real weight of each cause of maternal deaths around the world.

During the research, we realized that the majority of the diseases that appear during pregnancy are still understudied, the few scientific studies that have so far been allowed in pregnant women were not enough to understand the etiology, pathogenesis and even to test some therapeutic specifically in this stage of woman's life. Only in some diseases, like hypertension syndromes, hemorrhage and infection, there is a well-established prevention methodology. Surviving sepsis campaign, as others campaigns, had a good impact in death reduction in non-pregnant women due to establishing guidelines, standardized treatment and greater investment in training of health professionals. Perhaps it is time to apply this strategy to pregnant women.

In order to compare the situation of Portugal with other European countries, we researched their respective reports and we realize that, like Portugal they weren't updated for many years.^[58]^[59] A Swedish study shows a pregnancy related mortality ratio of 7.3 which is 64% highest than the report of World Health Organization.^[58] But just as Portugal, the Swedish report is from 2007.^[58]

The Spaniards report includes statistic numbers between 2001 and 2005 show a RMM of 4.1 and a increase in 20% of the number of pregnant woman with 35 years or over since 1996.^[59]

United Nations created, in 2000, the Millennium Development Goals (MDGs), which include maternal death surveillance and response (MDSR) in order to improve the precision of the notification of the deaths. The aim of MDSR is to get less than 30 deaths per 100000 pregnancies until 2030.^[1]

In each hour, 30 women die around the world from pregnancy complications. Some of maternal death can be avoided with better individual heath cares so it is very important to prioritize the health politics and funding programs to achieve better data on maternal mortality and thus improve the health indices.^[55]

Although it still has a long way to go, MRSD studies are the beginning for a new era with a more accurate view of this major problem that is maternal mortality. Review on this subject

allows saving the lives of many pregnant women and their newborns.

References:

1. Tara M Herrick, C.M.H.-J., Alice M Levisay, et al, *Prioritizing investments in innovations to protect women from the leading causes of maternal death*. BMC Pregnancy and Childbirth, 2014. **14**.
2. WHO, U., UNFPA, World Bank Group and the United Nations Population Division, *Trends in maternal mortality: 1990 to 2015 Estimates by WHO, UNICEF, UNFPA, World Bank Group and the United Nations Population Division*. 2015. p. 12.
3. Sharma, B.R.e.a., *Forensic considerations of pregnancy-related maternal deaths: An overview*. Journal of Forensic and Legal Medicine 2009. **16**(2009): p. 233-238.
4. Belizán, S.M.a.J.M., *The true cost of maternal death: individual tragedy impacts family, community and nations*, B. central, Editor. 2015.
5. *Principais Indicadores da Saúde para Portugal 2015*, DGS. p. 1.
6. S, L., *Maternal Death, Autopsy Studies, and Lessons from Pathology*. . PLoS Med, 2008. **2**(5): p. e48.
7. Roch Cantwell, T.C., Griselda Cooper, Andrew Dawson, James Drife MD, Debbie, Garrod RM, Ann Harper *Saving Mothers' Lives: Reviewing maternal deaths to make motherhood safer—2006–08*. Vol. 118. 2011. 205.
8. David Dolinak , E.M., Emma Lew, *Forensic Pathology: Principles and Practice*. 1st ed. 616.
9. Roger, M., *Pregnancy and venous thromboembolism: 'TIPPS' for risk stratification*. 2014, American Society of Hematology: Reproductive Reasoning: Hematology and Reproduction. p. 387-392.
10. Cavalcanti, E.F.d.A.M., Herlon Saraiva., *Clínica médica: dos sinais e sintomas ao diagnóstico e tratamento*. . 2007: p. 1958.
11. F. Cunningham, K.L., S. Bloom, et al, *Williams OBSTETRICS*. 24th ed. 2014.
12. Alexander M. Friedman, M.C.V.A., PhD et al, *Thromboembolism incidence and prophylaxis during vaginal delivery hospitalizations*. American Journal of Obstetrics & Gynecology, 2015. **221**: p. e1- e12.
13. Alexander M. Friedman, M.C.V.A., PhD *Obstetrical venous thromboembolism: Epidemiology and strategies for prophylaxis*. Elsevier, 2015. **0**(0).
14. Kenya Kamimura, H.A., Hirokazu Kawai et al, *Advances in understanding and treating liver diseases during pregnancy: A review*. World J Gastroenterol, 2015. **21**(17): p. 5183-5190.
15. Charles R. B. Beckmann, F.W.e.a., *Obstetrics and Gynecology*. 6ª ed. 497.
16. Graça, L.M.D., *Medicina Materbo-Fetal*. 4ª ed.
17. Yan Zhong, F.Z.a.Y.D., *Serum screening in first trimester to predict pre-eclampsia, small for gestational age and preterm delivery: systematic review and meta-analysis*. BMC Pregnancy and Childbirth, 2015. **15**: p. 191.
18. Alex R. Souza, M.R.A., Aurélio A. R. Costa, Carlos N. Neto, *Tratamento anti-hipertensivo na gravidez* Acta Med Portuguesa, 2010. **23**: p. 077-084.
19. Stout MJ1, C.S., Colditz GA, et al, *The Utility of 12-Hour Urine Collection for the Diagnosis of Preeclampsia: A Systematic Review and Meta-analysis*. Obstet Gynecol, 2015. **126**(4): p. 731-736.
20. Haram, K.S., E. Abildgaard, U., *The HELLP syndrome: clinical issues and management. A Review*. BMC Pregnancy Childbirth, 2009. **9**: p. 8.
21. De Oliveira, R., De Matos, IC., Da Silva, TBP et al, *Síndrome Hellp: estudo de revisão para o cuidado de enfermagem* Enfermaria global, 2012. **28**: p. 346-354.
22. Lisa R. Leffert, M.C.R.C., B et al, *Patient Characteristics and Outcomes After Hemorrhagic Stroke in Pregnancy*. Circulation Cardiovascular Quality and Outcomes, 2015. **8**(6): p. S170-178.
23. Florin-Andrei Taran, K.-O.K., Markus Hübner et al, *The Diagnosis and Treatment of Ectopic Pregnancy*. Deutsches Ärzteblatt International, 2015. **37**(4): p. 693-704.
24. Vyshka, G.C., N. Shaqiri, E., *Placenta increta causing hemoperitoneum in the 26th week of pregnancy: a case report*. J Med Case Rep, 2010. **4**: p. 412.

25. al, F.R.C.R.M.Y.N.T.R.S.M.e., *Fetal death in placental abruption: comparison of two different time periods*. Revista da Associação Médica Brasileira, 2008. **54**.
26. Theodoros Pavlis, C.S., Stavros Gourgiotis et al, *Spontaneous Rupture of Splenic Artery Aneurysm during the First Trimester of Pregnancy: Report of an Extremely Rare Case and Review of the Literature*. **Hindawi Publishing Corporation**, 2012. **Obstetrics and Gynecology**: p. 3.
27. Betal, D., et al., *Spontaneous ruptured splenic artery aneurysm: a case report*. Cases J, 2009. **2**: p. 7150.
28. Veluppillai C, P.S., de Kerviler B et al *Splenic arterial aneurysm and pregnancy: A review*. Presse Med, 2015. **44**(10): p. 991-994.
29. McDonnell, N.J., V. Percival, and M.J. Paech, *Amniotic fluid embolism: a leading cause of maternal death yet still a medical conundrum*. Int J Obstet Anesth, 2013. **22**(4): p. 329-36.
30. Balinger KJ, C.L.M., Hon HH, et al, *Amniotic fluid embolism: despite progress, challenges remain*. Curr Opin Obstet Gynecol, 2015. **27**(6): p. 398- 405.
31. Rosalinda Rodrigues, F.N., Manuel Meirinho *Embolia de liquido amniótico*. Acta Medica Portuguesa, 2007. **20**: p. 93-96.
32. Rita Torres, N.M., Sara Valadares et al, *Maternal sepsis*. Acta Obstet Ginecol Port, 2015. **9**(1): p. 65-72.
33. Buddeberg BS, A.W., *Puerperal sepsis in the 21st century: progress, new challenges and the situation worldwide*. Postgrad Med J, 2015. **91**: p. 572-578.
34. Ricardo Luiz Cordioli, E.C., Romulo Negrini, et al, *Sepsis and pregnancy: do we know how to treat this situation?* Rev Bras Ter Intensiva, 2013. **25**(4): p. 334-344.
35. Tsokos, M., *Pathology of Sepsis*, in *Essentials of Autopsy Practice*, G. Rutty, Editor. 2006, Springer London. p. 39-85.
36. James S. Ware, P.D., Jian Li, Ph.D., Erica Mazaika, et al, *Shared Genetic Predisposition in Peripartum and Dilated Cardiomyopathies*. The NEW ENGLAND JOURNAL of MEDICINE, 2016.
37. Ellawindy A, S.K., Sunamura S, et al, *Rho-Kinase Inhibition During Early Cardiac Development Causes Arrhythmogenic Right Ventricular Cardiomyopathy in Mice*. Arterioscler Thromb Vasc Biol, 2015. **35**(10).
38. Agir, A.B., S. Celikyurt, U. Argan, O. Yilmaz, I. Karauzum, K. Vural, A., *Arrhythmogenic right ventricular cardiomyopathy in pregnancy*. Int Heart J, 2014. **55**(4): p. 372-6.
39. Sebastien P.J. Krul, J.J.v.d.S., et al, *Systematic review of pregnancy in women with inherited cardiomyopathies*. European Journal of Heart Failure 2011. **13**: p. 584-594.
40. Allen Patrick Burke, M.C.E.A.P.B., MD, *Arrhythmogenic Right Ventricular Cardiomyopathy Pathology*. 2015: Medscape.
41. S., S., *Spontaneous rupture of acute ascending aortic dissection in a young pregnant woman: a sudden unexpected death*. Forensic Sci Int., 2013: p. e5-8.
42. Hind N. Moussa, M.M.M., MD Mateo G. Leon, MD et al, *Acute Myocardial Infarction Due to Coronary Artery Dissection in the Postpartum Period*. Am J Perinatol, 2015. **05**(02): p. e93-e96.
43. Samanta Soares, Â.F., Ivone Lobo, et al, *Acute myocardial infarction in pregnancy – case report*. Acta Obstet Ginecol Port 2010. **4**(4): p. 213-216.
44. Edey, S.M., N. Nashef, L., *SUDEP and epilepsy-related mortality in pregnancy*. Epilepsia, 2014. **55**(7): p. e72-4.
45. Ozhan Ozdemir, M.E.S., Aslihan Kurt,, *Pregnancy outcome of 149 pregnancies in women with epilepsy: Experience from a tertiary care hospital*. Interventional Medicine & Applied Science, 2015. **7**(3): p. 108-113.
46. *The American Pregnancy Association*. [cited 2015 28.11]; Available from: <http://americanpregnancy.org/about/>.
47. *Neurolis: clinica de neurologia e neurofisiologia. Lda*. [cited 2015 28.11]; Available from: <http://www.neurolis.pt/index.php/epilepsia/diagnostico-da-epilepsia>.
48. Shahn Nouri, M.C.E.S.R.B., MD *Sudden Unexpected Death in Epilepsy*. 2013: Medscape.
49. Yamamoto, T.F., Y. Emoto, Y. et al, *Autopsy case of sudden maternal death from thrombotic thrombocytopenic purpura*. J Obstet Gynaecol Res, 2013. **39**(1): p. 351-4.
50. Carpenter, M.R.a.S.L., *Thrombotic Microangiopathies*. ISRN Hematology, 2012. **2012**: p. 8.
51. Plu, I., et al., *Pregnancy, cesarean, and pheochromocytoma: a case report and literature review*. J Forensic Sci, 2013. **58**(4): p. 1075-9.

52. Cecchi R, F.P., Capri O, Cipolloni L., *A rare case of sudden death due to hypotension during cesarean section in a woman suffering from pheochromocytoma and neurofibromatosis*. J Forensic Sci., 2013. **6**: p. 1636-9.
53. J. G. Hudsmith, C.E.T., D. A. Browne, *Undiagnosed phaeochromocytoma mimicking severe preeclampsia in a pregnant woman at term*. International Journal of Obstetric Anesthesia, 2006. **15**: p. 240-254.
54. Gomes, M.T.V.M.d.C., *Mortes Maternas em Portugal, 2001 - 2007*. 2009, DGS. p. 32.
55. Steven L. Clark, M.M.A.B., et al, *Maternal death in the 21st century: causes, prevention, and relationship to cesarean delivery*. American Journal of Obstetrics & Gynecology, 2008: p. 36. e1- e5.
56. Tsokos, C.B.M.S.M., *Maternal and pregnancy-related death: causes and frequencies in an autopsy study population*. Forensic Sci Med Pathol, 2013. **9**(3): p. 296-307.
57. Lewis, G., *Reviewing maternal deaths to make pregnancy safer*. Best Practice & Research Clinical Obstetrics and Gynaecology, 2008. **22**: p. 447-463.
58. Esscher, A.H., U. Haglund, B. et al, *Maternal mortality in Sweden 1988-2007: more deaths than officially reported*. Acta Obstet Gynecol Scand, 2013. **92**(1): p. 40-6.
59. Luque Fernandez, M.A.C., A. B. Dramaix-Wilmet, M. et al, *Increase in maternal mortality associated with change in the reproductive pattern in Spain: 1996-2005*. J Epidemiol Community Health, 2009. **63**(6): p. 433-8.

ANO	MM	NV	RMM
1993	10	114030	8,8
1994	12	109287	11,0
1995	9	107184	8,41
1996	8	110363	7,2
1997	14	113047	12,4
1998	6	113510	5,3
1999	6	116038	5,2
2000	3	120071	2,5
2001	15	112825	13,3
2002	15	114456	13,1
2003	10	112589	8,9
2004	15	109356	13,7
2005	6	109457	5,5
2006	20	105514	19,0
2007	11	102567	10,7

Table 1. Number of maternal deaths (MM), live births (NV) and MMR between 1993 and 2007 in Portugal (adapted from Directorate-General of Health) ^[54]

AGE	2001	2002	2003	2004	2005	2006	2007	TOTAL	%
15-19				1	1	1	1	4	4,3
20-24	2	1	2	1				6	6,5
25-29	2	4	2	3		4	1	16	17,4
30-34	2	6	5	4	3	9	2	31	33,7
35-39	6	3		3	2	5	7	26	28,3
40-44	2	1	1	2		1		7	7,6
>44	1			1				2	2,2
TOTAL	15	15	10	15	6	20	11	92	100,0

Table 2- Maternal mortality by age group between 2001 and 2007 (adapted from Directorate-General of Health) ^[54]

Causes of maternal death	2001	2002	2003	2004	2005	2006	2007	TOTAL	% Of total of known causes
Direct								52	72,2
Haemorrhage/ coagulopathy	6	1	1	2	2	4	3	19	26,4
Hypertensive syndrome of pregnancy	3	4	2			7	1	17	23,6
Infection/ sepsis	1		1	3		1	1	7	9,7
Tromboembolis		1	1		3		1	6	8,3
Others embolis					1	1	1	3	4,2
Indirect								18	25,0
Infection		1	1		2		4		5,6
Cardiac death	2	1	1	1		1		6	8,3
Cancer	1	1						2	2,8
CNS		1				1		2	2,8
Others		1		1		2		4	5,6
Accidental/ incidental								2	2,8
Accidental/suicide				1			1		2,8
Complete with known causes	13	11	7	8	6	19	8	72	100,0
Unknown	2	4	3	7		1	3	20	21,7
TOTAL	15	15	10	15	6	20	11	92	

Table 3- Causes of maternal death between 2001 and 2007(adapted from Directorate-General of Health) ^[54]




Figure 1. Evolution of MMR, number of maternal deaths per 100,000 live births between 1975 and 2007(adapted from Directorate-General of Health) ^[54]

MDG region	MMR	Range of MMR uncertainty (80% UI)		Number of maternal deaths	Lifetime risk of maternal death 1 in:
		Lower	Upper		
World	216	207	249	303000	180
Developed regions	12	11	14	1700	4900
Developing regions	239	229	275	302000	150
Northern africa	70	56	92	3100	450
Sub-saharan africa	546	511	652	201000	36
Easten asia	27	23	33	4800	2300
Eastern asia excluding china	43	24	86	378	1500
Southern asia	176	153	216	66000	210
Southern asia excluding india	180	147	249	21000	190
South- eastern asia	110	95	142	13000	380
Western asia	91	73	125	4700	360
Caucasus and central asia	33	27	45	610	1100
Latin america and the caribbean	67	64	77	7300	670
Latin america	60	57	66	6600	760
Caribbean	175	130	265	1300	250
Oceania	187	95	381	500	150

Table 4- estimates of maternal mortality ratio(MMR, maternal deaths per 100000 live births), number of matenal deaths, and lifetime risk, by National Millennium Development Goal (MDG) region, 2015(adapted from WHO) [2]

Agradecimentos

A realização desta monografia contou com importantes apoios e incentivos sem os quais não se teria tornado uma realidade e aos quais estarei eternamente grata.

Ao Professor Doutor Agostinho Santos, pela sua orientação, paciência e disponibilidade, pelas opiniões e pelas críticas, pela total colaboração para solucionar problemas que foram surgindo ao longo da realização deste trabalho, permitindo assim elevar a qualidade e o impacto do trabalho.

À Doutora Joana Félix, pela disponibilização de material como livros e artigos sobre o tema e pela disponibilidade para rever alguns conceitos específicos da área de Obstetrícia.

Anexos



JOURNAL OF FORENSIC AND LEGAL MEDICINE

An International journal of Forensic and Legal Medicine

AUTHOR INFORMATION PACK

TABLE OF CONTENTS

•	Description	p.1
•	Audience	p.1
•	Impact Factor	p.1
•	Abstracting and Indexing	p.2
•	Editorial Board	p.2
•	Guide for Authors	p.4



ISSN: 1752-928X

DESCRIPTION

Official journal of the [Faculty of Legal and Forensic Medicine](#).

The *Journal of Forensic and Legal Medicine* publishes topical articles on legal medicine and all clinical aspects of forensic medicine and related specialities. All submissions are peer-reviewed by at least two independent reviewers, and the Journal is listed in MEDLINE/Index Medicus. All aspects of legal medicine, the coronial system and the medical principles of care and forensic assessment of living individuals, whether adult or child, in contact with the judicial system are examined and the Journal has a broad international perspective. Topics covered in the journal include, but are not necessarily limited to, the following, particularly with relevance to the clinical setting: forensic medicine, forensic pathology, forensic science, forensic toxicology, forensic odontology, forensic anthropology, forensic psychiatry, forensic genetics, death and care in custody, training, forensic nursing, occupational health of police, fitness to interview, death investigation and causes of death, forensic medicine, national systems, human identification, entomology and palynology, taphonomy, mental health, criminology, profiling, child abuse and neglect, interpersonal violence, assault and injury, personal injury, elder abuse, domestic violence, sexual assault, suicide, parasuicide and deliberate self-harm, restraint injuries, drug and alcohol misuse, drink and drug driving, traffic medicine, transportation medicine, refugee and asylum medicine, medical law, medical ethics, euthanasia and end of life issues, consent, confidentiality, clinical negligence, professional regulation, patients' complaints procedures, medical disciplinary procedures, the coronial system, clinical governance, clinical risk management, clinical performance review procedures, criminal charges arising from clinical practice such as murder, manslaughter and indecent assault, the sick doctor, the validation, licensing and certification of doctors, mass disaster, war graves, torture, extra-judicial death, human rights.

AUDIENCE

Police Surgeons, Forensic Clinicians, Scientists, Psychiatrists, Pathologists, Odontologists and Nurses. Also Coroners, Prison Medical Officers, Criminologists, Lawyers, Police Officers, ER Staff and Armed Forces Medical Officers

IMPACT FACTOR

2014: 0.760 © Thomson Reuters Journal Citation Reports 2015

ABSTRACTING AND INDEXING

Cambridge Scientific Abstracts
Criminal Justice Abstracts
MEDLINE®
EMBASE
National Criminal Justice Reference Service
Bibliography of Periodical Literature and International Bibliography of Book Reviews
Forensic Science FORS
Scopus

EDITORIAL BOARD

Editor-in-Chief

J Payne-James

Associate Editors

P. Chariot, Bondy Cedex, France
C.M. Milroy, Ottawa, Ontario, Canada

International Editorial Board

A. Aggrawal, New Delhi, India
D. Alempijevic, Belgrade, Serbia
W. Anderson, Reading, UK
P. Beh, Hong Kong, China
S. Black, Dundee, UK, Scotland
T. Bleetman, Northants, UK
J. Clark, Glasgow, UK, Scotland
N. Clements, Leeds, UK
D. Cusack, Dublin, Ireland
M. Devin, York, England, UK
J.V. Evans, Ilkley, UK
G. Fernie, Edinburgh, UK
J. Freedman, Suffolk, England, UK
J. Gall, Melbourne, VIC, Australia
P. Green, London, UK
G. Gudjonsson, London, UK
C. Hall, Victoria, BC, Canada
V. Iacopino, Henderson, Nevada, USA
A.W. Jones, Stockholm, Sweden
S.E. Josse, London, UK
K. Kennedy, Galway, Ireland
A. Keogh, Standish, UK
M. Kharoshah, Cairo, Egypt
N.P. Lemos, San Francisco, CA, USA
C. Lincoln, Queensland, Northern Territory, Australia
P. Marks, Leeds, UK
R.G. Menezes, Dammam, Saudi Arabia
R. Metcalf, Fort Worth, TX, USA
K. Nadesan, New Lambton, NSW, Australia
M. Odell, Victoria, VIC, Australia
O. Ozcalipsi, Istanbul, Turkey
D.J. Pounder, Dundee, UK, Scotland
D. Ranson, Southbank, VIC, Australia
P. Saukko, Turku, Finland
P. Schütte, Norfolk, England, UK
R.D. Sheridan, Salisbury, Wiltshire, UK, England
M. Stark, London, UK, England
T. Thompson, Middlesbrough, UK, England
J.L. Thomsen, Odense C, Denmark
K. Törő, Budapest, Hungary
R. Tunbridge, Crowthorne, UK
D.N. Vieira, Coimbra, Portugal
G. Vilke, San Diego, CA, USA
M. Welner, New York, New York, USA

C. White, Manchester, UK
A. Wilkinson, Oxford, England, UK
A. Williams, South Melbourne, VIC, Australia

GUIDE FOR AUTHORS

INTRODUCTION

Types of paper

The following types of articles will be considered for publication:

Original Communication: new research, previously unpublished.

Review article: detailed review of specific subject, backed up by full reference list and exploring all aspects of subject.

Clinical Practice: review backed up by relevant literature of specific aspects of clinical practice.

Short Report: new research or clinical issue, straightforward idea, simple methodology, concise take-home message.

Leading Article: invited article by an authority on a particular issue.

Editorial: topical polemic on an issue of the day, some commissioned, some submitted.

Letter to the Editor: comment or useful critique on material published in the journal. The decision to publish submitted letters rests purely with the Editor-in-Chief.

Book Review: review of relevant books which are not more than 2 years old. Unsolicited reviews will not usually be accepted, but suggestions for appropriate books for review should be sent to the Editor-in-Chief.

Consideration will be given by the Editor to other categories of article that do not fit into the above. Contact the Editor at jasonpaynejames@aol.com for any specific inquiries on these matters.

Contact details for submission

Authors should send queries concerning the submission process or journal procedures to AuthorSupport@elsevier.com. Authors can check the status of their manuscript within the review procedure using Elsevier Editorial System.

BEFORE YOU BEGIN

Ethics in publishing

For information on Ethics in publishing and Ethical guidelines for journal publication see <https://www.elsevier.com/publishingethics> and <https://www.elsevier.com/journal-authors/ethics>.

Human and animal rights

If the work involves the use of human subjects, the author should ensure that the work described has been carried out in accordance with The Code of Ethics of the World Medical Association (Declaration of Helsinki) for experiments involving humans, <http://www.wma.net/en/30publications/10policies/b3/index.html>; Uniform Requirements for manuscripts submitted to Biomedical journals, <http://www.icmje.org>. Authors should include a statement in the manuscript that informed consent was obtained for experimentation with human subjects. The privacy rights of human subjects must always be observed.

All animal experiments should be carried out in accordance with the U.K. Animals (Scientific Procedures) Act, 1986 and associated guidelines, [EU Directive 2010/63/EU for animal experiments](#), or the National Institutes of Health guide for the care and use of Laboratory animals (NIH Publications No. 8023, revised 1978) and the authors should clearly indicate in the manuscript that such guidelines have been followed. **All animal studies need to ensure they comply with the ARRIVE guidelines.** More information can be found at <http://www.nc3rs.org.uk/page.asp?id=1357>.

Conflict of interest

All authors must disclose any financial and personal relationships with other people or organizations that could inappropriately influence (bias) their work. Examples of potential conflicts of interest include employment, consultancies, stock ownership, honoraria, paid expert testimony, patent applications/registrations, and grants or other funding. If there are no conflicts of interest then please state this: 'Conflicts of interest: none'. See also <https://www.elsevier.com/conflictsofinterest>. Further information and an example of a Conflict of Interest form can be found at: http://service.elsevier.com/app/answers/detail/a_id/286/supporthub/publishing.

Submission declaration

Submission of an article implies that the work described has not been published previously (except in the form of an abstract or as part of a published lecture or academic thesis or as an electronic preprint, see <https://www.elsevier.com/sharingpolicy>), that it is not under consideration for publication elsewhere, that its publication is approved by all authors and tacitly or explicitly by the responsible authorities where the work was carried out, and that, if accepted, it will not be published elsewhere including electronically in the same form, in English or in any other language, without the written consent of the copyright-holder.

Authorship

All authors should have made substantial contributions to all of the following: (1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data, (2) drafting the article or revising it critically for important intellectual content, (3) final approval of the version to be submitted.

Changes to authorship

Authors are expected to consider carefully the list and order of authors **before** submitting their manuscript and provide the definitive list of authors at the time of the original submission. Any addition, deletion or rearrangement of author names in the authorship list should be made only **before** the manuscript has been accepted and only if approved by the journal Editor. To request such a change, the Editor must receive the following from the **corresponding author**: (a) the reason for the change in author list and (b) written confirmation (e-mail, letter) from all authors that they agree with the addition, removal or rearrangement. In the case of addition or removal of authors, this includes confirmation from the author being added or removed.

Only in exceptional circumstances will the Editor consider the addition, deletion or rearrangement of authors **after** the manuscript has been accepted. While the Editor considers the request, publication of the manuscript will be suspended. If the manuscript has already been published in an online issue, any requests approved by the Editor will result in a corrigendum.

Article transfer service

This journal is part of our Article Transfer Service. This means that if the Editor feels your article is more suitable in one of our other participating journals, then you may be asked to consider transferring the article to one of those. If you agree, your article will be transferred automatically on your behalf with no need to reformat. Please note that your article will be reviewed again by the new journal. More information about this can be found here: <https://www.elsevier.com/authors/article-transfer-service>.

Unpublished material

Material in unpublished letters and manuscripts is also protected and must not be published unless permission has been obtained.

Copyright

Upon acceptance of an article, authors will be asked to complete a 'Journal Publishing Agreement' (for more information on this and copyright, see <https://www.elsevier.com/copyright>). An e-mail will be sent to the corresponding author confirming receipt of the manuscript together with a 'Journal Publishing Agreement' form or a link to the online version of this agreement.

Subscribers may reproduce tables of contents or prepare lists of articles including abstracts for internal circulation within their institutions. Permission of the Publisher is required for resale or distribution outside the institution and for all other derivative works, including compilations and translations (please consult <https://www.elsevier.com/permissions>). If excerpts from other copyrighted works are included, the author(s) must obtain written permission from the copyright owners and credit the source(s) in the article. Elsevier has preprinted forms for use by authors in these cases: please consult <https://www.elsevier.com/permissions>.

For open access articles: Upon acceptance of an article, authors will be asked to complete an 'Exclusive License Agreement' (for more information see <https://www.elsevier.com/OAauthoragreement>). Permitted third party reuse of open access articles is determined by the author's choice of user license (see <https://www.elsevier.com/openaccesslicenses>).

Author rights

As an author you (or your employer or institution) have certain rights to reuse your work. For more information see <https://www.elsevier.com/copyright>.

Role of the funding source

You are requested to identify who provided financial support for the conduct of the research and/or preparation of the article and to briefly describe the role of the sponsor(s), if any, in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the article for publication. If the funding source(s) had no such involvement then this should be stated.

Funding body agreements and policies

Elsevier has established a number of agreements with funding bodies which allow authors to comply with their funder's open access policies. Some authors may also be reimbursed for associated publication fees. To learn more about existing agreements please visit <https://www.elsevier.com/fundingbodies>.

After acceptance, open access papers will be published under a noncommercial license. For authors requiring a commercial CC BY license, you can apply after your manuscript is accepted for publication.

Open access

This journal offers authors a choice in publishing their research:

Open access

- Articles are freely available to both subscribers and the wider public with permitted reuse
- An open access publication fee is payable by authors or on their behalf e.g. by their research funder or institution

Subscription

- Articles are made available to subscribers as well as developing countries and patient groups through our universal access programs (<https://www.elsevier.com/access>).
- No open access publication fee payable by authors.

Regardless of how you choose to publish your article, the journal will apply the same peer review criteria and acceptance standards.

For open access articles, permitted third party (re)use is defined by the following Creative Commons user licenses:

Creative Commons Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

For non-commercial purposes, lets others distribute and copy the article, and to include in a collective work (such as an anthology), as long as they credit the author(s) and provided they do not alter or modify the article.

The open access publication fee for this journal is **USD 3000**, excluding taxes. Learn more about Elsevier's pricing policy: <http://www.elsevier.com/openaccesspricing>.

Green open access

Authors can share their research in a variety of different ways and Elsevier has a number of green open access options available. We recommend authors see our green open access page for further information (<http://elsevier.com/greenopenaccess>). Authors can also self-archive their manuscripts immediately and enable public access from their institution's repository after an embargo period. This is the version that has been accepted for publication and which typically includes author-incorporated changes suggested during submission, peer review and in editor-author communications. Embargo period: For subscription articles, an appropriate amount of time is needed for journals to deliver value to subscribing customers before an article becomes freely available to the public. This is the embargo period and it begins from the date the article is formally published online in its final and fully citable form.

This journal has an embargo period of 12 months.

Language (usage and editing services)

Please write your text in good English (American or British usage is accepted, but not a mixture of these). Authors who feel their English language manuscript may require editing to eliminate possible grammatical or spelling errors and to conform to correct scientific English may wish to use the English Language Editing service available from Elsevier's WebShop (<http://webshop.elsevier.com/languageediting/>) or visit our customer support site (<http://support.elsevier.com>) for more information.

Submission

Our online submission system guides you stepwise through the process of entering your article details and uploading your files. The system converts your article files to a single PDF file used in the peer-review process. Editable files (e.g., Word, LaTeX) are required to typeset your article for final publication. All correspondence, including notification of the Editor's decision and requests for revision, is sent by e-mail.

For certain submissions, authors should supply the names and email addresses of three potential reviewers for their manuscript, all of which must be in a different country. Please do not suggest reviewers from your own institution, previous or current collaborators, or Editorial Board members. Also, please do not contact any reviewers you have suggested.

Submit your article

Please submit your article via <http://ees.elsevier.com/jflm>

Review Policy

As this journal has adopted a double blind reviewing policy, **please remove all identifying features** from the paper itself by ensuring that no author's name appears in the main text, in-text citations, reference list, or any running header. Author information should *only* be included in the title page.

PREPARATION

Subdivision - unnumbered sections

Divide your article into clearly defined sections. Each subsection is given a brief heading. Each heading should appear on its own separate line. Subsections should be used as much as possible when cross-referencing text: refer to the subsection by heading as opposed to simply 'the text'.

Headings for experimental papers should follow the usual conventions: Introduction, Methods, Results, Discussion, Acknowledgments. Other papers may be subdivided as the authors desire. The use of headings enhances readability.

Essential title page information

- **Title.** Concise and informative. Titles are often used in information-retrieval systems. Avoid abbreviations and formulae where possible.
- **Author names and affiliations.** Please clearly indicate the given name(s) and family name(s) of each author and check that all names are accurately spelled. Present the authors' affiliation addresses (where the actual work was done) below the names. Indicate all affiliations with a lower-case superscript letter immediately after the author's name and in front of the appropriate address. Provide the full postal address of each affiliation, including the country name and, if available, the e-mail address of each author.
- **Corresponding author.** Clearly indicate who will handle correspondence at all stages of refereeing and publication, also post-publication. **Ensure that the e-mail address is given and that contact details are kept up to date by the corresponding author.**
- **Present/permanent address.** If an author has moved since the work described in the article was done, or was visiting at the time, a 'Present address' (or 'Permanent address') may be indicated as a footnote to that author's name. The address at which the author actually did the work must be retained as the main, affiliation address. Superscript Arabic numerals are used for such footnotes.

Abstract

A concise and factual abstract is required. The abstract should state briefly the purpose of the research, the principal results and major conclusions. An abstract is often presented separately from the article, so it must be able to stand alone. For this reason, References should be avoided, but if essential, then cite the author(s) and year(s). Also, non-standard or uncommon abbreviations should be avoided, but if essential they must be defined at their first mention in the abstract itself.

Graphical abstract

Although a graphical abstract is optional, its use is encouraged as it draws more attention to the online article. The graphical abstract should summarize the contents of the article in a concise, pictorial form designed to capture the attention of a wide readership. Graphical abstracts should be submitted as a separate file in the online submission system. Image size: Please provide an image with a minimum of 531 × 1328 pixels (h × w) or proportionally more. The image should be readable at a size of 5 × 13 cm using a regular screen resolution of 96 dpi. Preferred file types: TIFF, EPS, PDF or MS Office files. See <https://www.elsevier.com/graphicalabstracts> for examples.

Authors can make use of Elsevier's Illustration and Enhancement service to ensure the best presentation of their images and in accordance with all technical requirements: [Illustration Service](#).

Highlights

Highlights are mandatory for this journal. They consist of a short collection of bullet points that convey the core findings of the article and should be submitted in a separate editable file in the online submission system. Please use 'Highlights' in the file name and include 3 to 5 bullet points (maximum 85 characters, including spaces, per bullet point). See <https://www.elsevier.com/highlights> for examples.

Keywords

Immediately after the abstract, provide a maximum of 6 keywords, using American spelling and avoiding general and plural terms and multiple concepts (avoid, for example, 'and', 'of'). Be sparing with abbreviations: only abbreviations firmly established in the field may be eligible. These keywords will be used for indexing purposes.

Abbreviations

Avoid abbreviations in the title and abstract. All unusual abbreviations should be fully explained at their first occurrence in the text.

Acknowledgements

Collate acknowledgements in a separate section at the end of the article before the references and do not, therefore, include them on the title page, as a footnote to the title or otherwise. List here those individuals who provided help during the research (e.g., providing language help, writing assistance or proof reading the article, etc.).

Nomenclature and units

Proprietary names of drugs, instruments etc. should be indicated by the use of initial capital letters. All measurements should be expressed in SI or SI-derived units.

Artwork

Electronic artwork

General points

- Make sure you use uniform lettering and sizing of your original artwork.
- Embed the used fonts if the application provides that option.
- Aim to use the following fonts in your illustrations: Arial, Courier, Times New Roman, Symbol, or use fonts that look similar.
- Number the illustrations according to their sequence in the text.
- Use a logical naming convention for your artwork files.
- Provide captions to illustrations separately.
- Size the illustrations close to the desired dimensions of the published version.
- Submit each illustration as a separate file.

A detailed guide on electronic artwork is available on our website:

<https://www.elsevier.com/artworkinstructions>.

You are urged to visit this site; some excerpts from the detailed information are given here.

Formats

If your electronic artwork is created in a Microsoft Office application (Word, PowerPoint, Excel) then please supply 'as is' in the native document format.

Regardless of the application used other than Microsoft Office, when your electronic artwork is finalized, please 'Save as' or convert the images to one of the following formats (note the resolution requirements for line drawings, halftones, and line/halftone combinations given below):

EPS (or PDF): Vector drawings, embed all used fonts.

TIFF (or JPEG): Color or grayscale photographs (halftones), keep to a minimum of 300 dpi.

TIFF (or JPEG): Bitmapped (pure black & white pixels) line drawings, keep to a minimum of 1000 dpi.

TIFF (or JPEG): Combinations bitmapped line/half-tone (color or grayscale), keep to a minimum of 500 dpi.

Please do not:

- Supply files that are optimized for screen use (e.g., GIF, BMP, PICT, WPG); these typically have a low number of pixels and limited set of colors;
- Supply files that are too low in resolution;
- Submit graphics that are disproportionately large for the content.

Where illustrations must include recognisable individuals, living or dead and of whatever age, great care must be taken to ensure that consent for publication has been given. It is the authors' responsibility to obtain written permission to reproduce borrowed material (illustrations and tables) from the original publishers and authors.

Color artwork

Please make sure that artwork files are in an acceptable format (TIFF (or JPEG), EPS (or PDF), or MS Office files) and with the correct resolution. If, together with your accepted article, you submit usable color figures then Elsevier will ensure, at no additional charge, that these figures will appear in color online (e.g., ScienceDirect and other sites) regardless of whether or not these illustrations are reproduced in color in the printed version. **For color reproduction in print, you will receive information regarding the costs from Elsevier after receipt of your accepted article.** Please indicate your preference for color: in print or online only. For further information on the preparation of electronic artwork, please see <https://www.elsevier.com/artworkinstructions>.

Illustration services

Elsevier's WebShop (<http://webshop.elsevier.com/illustrationservices>) offers Illustration Services to authors preparing to submit a manuscript but concerned about the quality of the images accompanying their article. Elsevier's expert illustrators can produce scientific, technical and medical-style images, as well as a full range of charts, tables and graphs. Image 'polishing' is also available, where our illustrators take your image(s) and improve them to a professional standard. Please visit the website to find out more.

Tables

Please submit tables as editable text and not as images. Tables can be placed either next to the relevant text in the article, or on separate page(s) at the end. Number tables consecutively in accordance with their appearance in the text and place any table notes below the table body. Be sparing in the use of tables and ensure that the data presented in them do not duplicate results described elsewhere in the article. Please avoid using vertical rules.

Web references

Refrain from using *online references* if possible. When referring to internet sources, for example Wikipedia, please state so clearly, and indicate if this information can be checked and on which date you visited this online source.

Reference management software

Most Elsevier journals have their reference template available in many of the most popular reference management software products. These include all products that support Citation Style Language styles (<http://citationstyles.org>), such as Mendeley (<http://www.mendeley.com/features/reference-manager>) and Zotero (<https://www.zotero.org/>), as well as EndNote (<http://endnote.com/downloads/styles>). Using the word processor plug-ins from these products, authors only need to select the appropriate journal template when preparing their article, after which citations and bibliographies will be automatically formatted in the journal's style. If no template is yet available for this journal, please follow the format of the sample references and citations as shown in this Guide.

Users of Mendeley Desktop can easily install the reference style for this journal by clicking the following link:

<http://open.mendeley.com/use-citation-style/journal-of-forensic-and-legal-medicine>

When preparing your manuscript, you will then be able to select this style using the Mendeley plug-ins for Microsoft Word or LibreOffice.

Reference style

Text: Indicate references by superscript numbers in the text. The actual authors can be referred to, but the reference number(s) must always be given.

List: Number the references in the list in the order in which they appear in the text.

Examples:

Reference to a journal publication:

1. Van der Geer J, Hanraads JAJ, Lupton RA. The art of writing a scientific article. *J Sci Commun* 2010;**163**:51–9.

Reference to a book:

2. Strunk Jr W, White EB. *The elements of style*. 4th ed. New York: Longman; 2000.

Reference to a chapter in an edited book:

3. Mettam GR, Adams LB. How to prepare an electronic version of your article. In: Jones BS, Smith RZ, editors. *Introduction to the electronic age*, New York: E-Publishing Inc; 2009, p. 281–304. Note shortened form for last page number. e.g., 51–9, and that for more than 6 authors the first 6 should be listed followed by 'et al.' For further details you are referred to 'Uniform Requirements for Manuscripts submitted to Biomedical Journals' (J Am Med Assoc 1997;**277**:927–34) (see also http://www.nlm.nih.gov/bsd/uniform_requirements.html).

Video data

Elsevier accepts video material and animation sequences to support and enhance your scientific research. Authors who have video or animation files that they wish to submit with their article are strongly encouraged to include links to these within the body of the article. This can be done in the same way as a figure or table by referring to the video or animation content and noting in the body text where it should be placed. All submitted files should be properly labeled so that they directly relate to the video file's content. In order to ensure that your video or animation material is directly usable, please provide the files in one of our recommended file formats with a preferred maximum size of 150 MB. Video and animation files supplied will be published online in the electronic version of your article in Elsevier Web products, including ScienceDirect: <http://www.sciencedirect.com>. Please supply 'stills' with your files: you can choose any frame from the video or animation or make a separate image. These will be used instead of standard icons and will personalize the link to your video data. For more detailed instructions please visit our video instruction pages at <https://www.elsevier.com/artworkinstructions>. Note: since video and animation cannot be embedded in the print version of the journal, please provide text for both the electronic and the print version for the portions of the article that refer to this content.

AudioSlides

The journal encourages authors to create an AudioSlides presentation with their published article. AudioSlides are brief, webinar-style presentations that are shown next to the online article on ScienceDirect. This gives authors the opportunity to summarize their research in their own words and to help readers understand what the paper is about. More information and examples are available at <https://www.elsevier.com/audioslides>. Authors of this journal will automatically receive an invitation e-mail to create an AudioSlides presentation after acceptance of their paper.

Supplementary material

Supplementary material can support and enhance your scientific research. Supplementary files offer the author additional possibilities to publish supporting applications, high-resolution images, background datasets, sound clips and more. Please note that such items are published online exactly as they are submitted; there is no typesetting involved (supplementary data supplied as an Excel file or as a PowerPoint slide will appear as such online). Please submit the material together with the article and supply a concise and descriptive caption for each file. If you wish to make any changes to supplementary data during any stage of the process, then please make sure to provide an updated file, and do not annotate any corrections on a previous version. Please also make sure to switch off the 'Track Changes' option in any Microsoft Office files as these will appear in the published supplementary file(s). For more detailed instructions please visit our artwork instruction pages at <https://www.elsevier.com/artworkinstructions>.

Submission checklist

The following list will be useful during the final checking of an article prior to sending it to the journal for review. Please consult this Guide for Authors for further details of any item.

Ensure that the following items are present:

One author has been designated as the corresponding author with contact details:

- E-mail address
- Full postal address

All necessary files have been uploaded, and contain:

- Keywords
- All figure captions
- All tables (including title, description, footnotes)

Further considerations

- Manuscript has been 'spell-checked' and 'grammar-checked'
- References are in the correct format for this journal
- All references mentioned in the Reference list are cited in the text, and vice versa
- Permission has been obtained for use of copyrighted material from other sources (including the Internet)

Printed version of figures (if applicable) in color or black-and-white

- Indicate clearly whether or not color or black-and-white in print is required.

For any further information please visit our customer support site at <http://support.elsevier.com>.

Submission of figures, tables and captions

Please note: Figures and their corresponding captions can be submitted together in a single file within EES. Tables and their corresponding captions can also be submitted together in a single file within EES. Please select the appropriate option(s) in the submission items drop down list.

AFTER ACCEPTANCE

Availability of accepted article

This journal makes articles available online as soon as possible after acceptance. This concerns the accepted article (both in HTML and PDF format), which has not yet been copyedited, typeset or proofread. A Digital Object Identifier (DOI) is allocated, thereby making it fully citable and searchable by title, author name(s) and the full text. The article's PDF also carries a disclaimer stating that it is an unedited article. Subsequent production stages will simply replace this version.

Use of the Digital Object Identifier

The Digital Object Identifier (DOI) may be used to cite and link to electronic documents. The DOI consists of a unique alpha-numeric character string which is assigned to a document by the publisher upon the initial electronic publication. The assigned DOI never changes. Therefore, it is an ideal medium for citing a document, particularly 'Articles in press' because they have not yet received their full bibliographic information. Example of a correctly given DOI (in URL format; here an article in the journal *Physics Letters B*):

<http://dx.doi.org/10.1016/j.physletb.2010.09.059>

When you use a DOI to create links to documents on the web, the DOIs are guaranteed never to change.

Online proof correction

Corresponding authors will receive an e-mail with a link to our online proofing system, allowing annotation and correction of proofs online. The environment is similar to MS Word: in addition to editing text, you can also comment on figures/tables and answer questions from the Copy Editor. Web-based proofing provides a faster and less error-prone process by allowing you to directly type your corrections, eliminating the potential introduction of errors.

If preferred, you can still choose to annotate and upload your edits on the PDF version. All instructions for proofing will be given in the e-mail we send to authors, including alternative methods to the online version and PDF.

We will do everything possible to get your article published quickly and accurately. Please use this proof only for checking the typesetting, editing, completeness and correctness of the text, tables and figures. Significant changes to the article as accepted for publication will only be considered at this stage with permission from the Editor. It is important to ensure that all corrections are sent back to us in one communication. Please check carefully before replying, as inclusion of any subsequent corrections cannot be guaranteed. Proofreading is solely your responsibility.

Offprints

The corresponding author, at no cost, will be provided with 25 free paper offprints, or, alternatively, a personalized link providing 50 days free access to the final published version of the article on [ScienceDirect](http://www.sciencedirect.com). This link can also be used for sharing via email and social networks. For an extra charge, more paper offprints can be ordered via the offprint order form which is sent once the article is accepted for publication. Both corresponding and co-authors may order offprints at any time via Elsevier's WebShop (<http://webshop.elsevier.com/myarticleservices/offprints>). Authors requiring printed copies of multiple articles may use Elsevier WebShop's 'Create Your Own Book' service to collate multiple articles within a single cover (<http://webshop.elsevier.com/myarticleservices/booklets>).

Author orders

When your article is published, you can commemorate your publication with printed author copies of the journal issue, customized full-color posters, extra offprints, and more. Please visit <http://webshop.elsevier.com> to learn more.

AUTHOR INQUIRIES

You can track your submitted article at <https://www.elsevier.com/track-submission>. You can track your accepted article at <https://www.elsevier.com/trackarticle>. You are also welcome to contact Customer Support via <http://support.elsevier.com>.

© Copyright 2014 Elsevier | <http://www.elsevier.com>